

Modified UW-Madison Lymphoma Protocol for Dogs

Body wt (kg) _____
 M^2 _____

Induction (page 1):

		<u>DATE</u>	<u>DOSE</u>	<u>WT/M²</u>
Week 1	Day 1: Vincristine, 0.5-0.7 mg/m ² IV Prednisone, 2 mg/kg PO SID	_____ _____	_____ _____	_____
Week 2	Cyclophosphamide, 250 mg/m ² IV or PO Prednisone, 1.5 mg/kg PO SID	_____ _____	_____ _____	_____
Week 3	Vincristine, 0.5-0.7 mg/m ² IV Prednisone, 1 mg/kg PO SID	_____ _____	_____ _____	_____
Week 4	Doxorubicin, 30 mg/m ² IV Prednisone, 0.5 mg/kg PO SID	_____ _____	_____ _____	_____
Week 5	no injectable drugs, discontinue prednisone			
Week 6	Vincristine, 0.5-0.7 mg/m ² IV	_____	_____	_____
Week 7	Cyclophosphamide, 250 mg/m ² IV or PO	_____	_____	_____
Week 8	Vincristine, 0.5-0.7 mg/m ² IV	_____	_____	_____
Week 9	Doxorubicin, 30 mg/m ² IV	_____	_____	_____

Skip 1 week.

If patient is in complete remission at week 9, continue with treatments at 1 week intervals on week 11.

A CBC must be performed within 48 hours prior to chemotherapy administration.

Delay chemotherapy by one week if:

Neutrophil count \leq 1500 cells/ μ l

Platelet count \leq 50,000 cells/ μ l

Patient is exhibiting GI signs secondary to previous chemotherapy

If hemorrhagic cystitis occurs, discontinue cyclophosphamide and substitute in chlorambucil @ 1.4 mg/kg PO.

Modified UW-Madison Lymphoma Protocol for Dogs

Body wt (kg) _____
 M^2 _____

Maintenance (page 2):

		<u>DATE</u>	<u>DOSE</u>	<u>WT/M^2</u>
Week 11	Vincristine, 0.5-0.7 mg/ m^2 IV	_____	_____	_____
Week 12	Cyclophosphamide, 250 mg/ m^2 IV or PO	_____	_____	_____
Week 13	Vincristine, 0.5-0.7 mg/ m^2 IV	_____	_____	_____
Week 14	Doxorubicin, 30 mg/ m^2 IV	_____	_____	_____
Week 16	Vincristine, 0.5-0.7 mg/ m^2 IV	_____	_____	_____
Week 17	Cyclophosphamide, 250 mg/ m^2 IV or PO	_____	_____	_____
Week 18	Vincristine, 0.5-0.7 mg/ m^2 IV	_____	_____	_____
Week 19	Doxorubicin, 30 mg/ m^2 IV	_____	_____	_____

Total doxorubicin dose to date: _____mg

Total dose of doxorubicin in m^2 : _____

A CBC must be performed within 48 hours prior to chemotherapy administration.

Delay chemotherapy by one week if:

Neutrophil count \leq 1500 cells/ μ l

Platelet count \leq 50,000 cells/ μ l

Patient is exhibiting GI signs secondary to previous chemotherapy

If patient is in complete remission at week 19, discontinue all therapy and reevaluate monthly with a thorough physical examination.